



Science & Cycling

29-30 June 2016, Caen, France

Short term parasympathetic reactivation after the LSCT: a practical tool to predict and monitor cycling performance

Lamberts R.P. ^{1,2}

¹ Division of Orthopaedic Surgery, Stellenbosch University, South Africa

² Division of Exercise Science and Sports Medicine, , University of Cape Town, South Africa.

Keywords: Cycling, LSCT, Monitoring, HRR, RMSSD

Introduction

The measurement of heart rate recovery (HRR) and parasympathetic reactivation after exercise to predict and monitor changes in training status has gained popularity over the last few years (Daanen et al, 2012, Bellenger et al., 2016). More recently, short term parasympathetic reactivation (LnRMSSD) has gained popularity as a marker of training status. However it has not yet been established over which time frame short term LnRMSSD should be measured and if gender effects LnRMSSD measurement. Therefore the aim of this study was to determine the LnRMSSD time interval with the highest reliability, determine the relationship between LnRMSSD and cycling performance and, determine if LnRMSSD is similar in male cyclists

Methods

Fifty male (5.1 ± 0.6 W/kg) and twenty female (4.8 ± 0.3 W/kg) cyclists participated on the study. Both HRR and LnRMSSD were captured as part of the LSCT (Lamberts et al., 2014), which was used as a warm-up before a Peak Power Output (PPO) test (including VO_{2max} determination), and two days later before a 40km time trial (40km TT) test. The reliability of the measurement was determined over 6 LSCT's, while 30, 60 and 90 second LnRMSSD data intervals were captured. Relationships were analysed between the most reliable LnRMSSD measurement and PPO, VO_{2max} and 40km TT performance

Results

The reliability and associated typical error of the measurement (TEM) of the parasympathetic measurements were the lowest over the 60 second period (LnRMSSD_{60s}) (ICC: 0.99; TEM: 7.7%). The relationship between HRR and training status was weaker than the relationship LnRMSSD_{60s} and training status. Strong relationships were found between PPO & LnRMSSD_{60s} in male ($r = 0.93$; $p < 0.0001$) and female ($r = 0.85$; $p < 0.0001$) cyclists. Slightly weaker relationships were found between VO_{2max} & LnRMSSD_{60s} in male ($r = 0.71$; $p < 0.0001$) and female ($r = 0.63$; $p < 0.0001$) cyclists and 40km TT time & LnRMSSD_{60s} in male ($r = -0.83$; $p <$

0.0001) and female ($r = -0.63$; $p = 0.003$) cyclists. Although similar slopes were found in male and female cyclists, the y-axis intercept was significantly different in all three performance parameters relationships with LnRMSSD_{60s} ($p < 0.0001$).

Discussion

Short term parasympathetic reactivation after the LSCT is a promising new parameter that will assist in more accurately predict and monitor training status in cyclists. The most reliable LnRMSSD measurement was found over a 60s period (LnRMSSD_{60s}). Good relationships between LnRMSSD_{60s} and PPO, $\text{VO}_{2\text{max}}$ and 40km TT were found both in male and female cyclists. Although similar relationships were found in male and female cyclists, different y-axis intercepts were found. This finding indicates that LnRMSSD_{60s} values in female cyclists relate differently to training status than in male cyclists. Although this study used the relationship between LnRMSSD_{60s} and cycling performance parameters (PPO, $\text{VO}_{2\text{max}}$ and 40km TT time), the author would like to emphasize that the actual LnRMSSD_{60s} measurement should be used for monitoring purposes and not the predicted cycling performance values. In addition, the author would like to emphasize that a multi-factorial approach (e.g. power, cadence and RPE from the LSCT) should be used to predict, monitor, and fine tune training prescription in cycling.

References

- Bellenger CR et al. (2016) Sports Med. Feb 18 – Epub ahead of print
Daanen HAM et al. (2012). Int J of Sports Physiol and Performance. 7:251-260
Lamberts RP (2014) Int J Sports Physiol and Performance 9:610-614